

## Saving teeth from the brink: a call to rethink the management of periodontally compromised teeth

 **Gabriel Oliveira Cavezzi<sup>1</sup>**,  
 **Orlando Cavezzi Junior<sup>2</sup>**

<sup>1</sup> Universidade Estadual de Londrina, Faculdade de Odontologia. Paraná, Brazil.

<sup>2</sup> Universidade São Francisco - Bragança Paulista, Faculdade de Odontologia. São Paulo, Brazil.

Dear editor:

Teeth, designed to last a lifetime, may be lost for various reasons such as root fractures or extensive caries. However, teeth compromised by periodontal disease, with uncertain or questionable prognoses, often receive little attention and are prematurely given a “death sentence.” Meanwhile, the introduction of dental implants as a therapeutic alternative has transformed daily clinical practice, substantially improving oral health and patients’ quality of life. In this context, the high success rates of osseointegrated implants, supported by solid scientific evidence, have encouraged clinicians to pursue an optimal therapeutic model.

Nevertheless, this approach should be adopted with caution, as implants are not exempt from failures or complications. The indiscriminate placement of these prostheses, often associated with iatrogenesis and infections, creates major clinical challenges. Deciding which tooth to extract is far from trivial; it requires careful, evidence-based judgment. Initially, many teeth show improvement after periodontal treatment, while others do not, demanding more cautious decision-making. Therefore, in cases of uncertainty, preserving the tooth should be the preferred option. This conservative stance is evidence-based, as scientific literature has confirmed the longevity of periodontally treated and restored teeth (1, 2).

It is concerning that evidence-based knowledge regarding the management of teeth affected by periodontal disease is being underestimated, leading to an evident lack of effort to preserve them (3). Studies have shown that the lower the clinician’s level of training, the more frequently extraction is recommended, which leads to significant changes in the approach to patient care (4, 5). Some clinicians argue that periodontally compromised teeth should be extracted to preserve bone volume for future implants; however, this aggressive approach is not supported by current evidence (6). Therefore, replacing compromised teeth with implants should not be regarded as a definitive solution for periodontal disease.

**Received:** June 24, 2024

**Accepted:** January 12, 2025

**Online:** June 30, 2025



Open access article

© The authors

### Cite as:

Cavezzi G, Cavezzi Junior O. Saving teeth from the brink: a call to rethink the management of periodontally compromised teeth. *Rev Estomatol Heredia*. 2025; 35(2): 157-158. DOI: 10.20453/reh.v35i2.5521

Clinicians frequently face the dilemma of whether to retain teeth affected by periodontal disease, whether due to limited training, unawareness of the evidence, or even negligence. Considering only tooth survival rates is insufficient, since prognosis depends on the extent and severity of the disease. Teeth with severe periodontal damage can be treated and preserved through scaling and root planing, along with periodontal maintenance therapy. The literature shows that these procedures can provide abutment teeth for fixed prostheses with a good long-term prognosis (7). The presence of furcation involvement further complicates the decision to preserve the tooth, with extraction often being the most common choice despite evidence supporting favorable long-term outcomes (2, 8).

Currently, the growing demand for esthetic treatments and the strong influence from the dental industry have fostered a global tendency to recommend dental implants, even in cases where conservative management could be a feasible alternative. This market-driven dynamic —often guided more by commercial than scientific considerations—has significantly influenced clinical decision-making, decreasing the frequency of dental tissue-preserving treatments. Moreover, although implant therapy shows encouraging survival rates, complications are common, prompting reflection on whether periodontally compromised teeth might actually have a longer lifespan than implants (9, 10). In this regard, dental implants should be viewed as a therapeutic option reserved for cases in which tooth preservation is no longer possible, rather than as a routine substitute for the natural dentition.

#### Corresponding author:

Orlando Cavezzi Junior

✉ [cavezzi@gmail.com](mailto:cavezzi@gmail.com)

## REFERENCES

1. Hirschfeld L, Wasserman B. A long-term survey of tooth loss in 600 treated periodontal patients. *J Periodontol* [Internet]. 1978; 49(5): 225-237. Available from: <https://doi.org/10.1902/jop.1978.49.5.225>
2. De Beule F, Alsaadi G, Perić M, Brex M. Periodontal treatment and maintenance of molars affected with severe periodontitis (DPSI = 4): an up to 27-year retrospective study in a private practice. *Quintessence Int* [Internet]. 2017; 48(5): 391-405. Available from: <https://doi.org/10.3290/j.qi.a38059>
3. Lindhe J, Pacey L. There is an overuse of implants in the world and an underuse of teeth as targets for treatment [entrevista]. *Br Dent J* [Internet]. 2014; 217(8): 396-397. Available from: <https://doi.org/10.1038/sj.bdj.2014.930>
4. Persson GR, Attström R, Lang NP, Page RC. Perceived risk of deteriorating periodontal conditions. *J Clin Periodontol* [Internet]. 2003; 30(11): 982-989. Available from: <https://doi.org/10.1034/j.1600-051x.2003.00415.x>
5. Carcamo-España V, Cuesta N, Flores P, Chimenos-Küstner E, Estrugo A, López-López J. Compromised teeth preserve or extract: a review of the literature. *J Clin Med* [Internet]. 2022; 11(18): 5301. Available from: <https://doi.org/10.3390/jcm11185301>
6. Gotfredsen K, Carlsson GE, Jokstad A, Arvidson Fyrberg K, Berge M, Bergendal B, et al. Implants and/or teeth: consensus statements and recommendations. *J Oral Rehabil* [Internet]. 2008; 35(suppl 1): 2-8. Available from: <https://doi.org/10.1111/j.1365-2842.2007.01833.x>
7. Lulic M, Brägger U, Lang NP, Zwahlen M, Salvi GE. Ante's (1926) law revisited: a systematic review on survival rates and complications of fixed dental prostheses (FDPs) on severely reduced periodontal tissue support. *Clin Oral Implants Res* [Internet]. 2007; 18(S3): 63-72. Available from: <https://doi.org/10.1111/j.1600-0501.2007.01438.x>
8. Nibali L, Zavattini A, Nagata K, Di Iorio A, Lin GH, Needleman I, et al. Tooth loss in molars with and without furcation involvement—a systematic review and meta-analysis. *J Clin Periodontol* [Internet]. 2016; 43(2): 156-166. Available from: <https://doi.org/10.1111/jcpe.12497>
9. Sartoretto SC, Shibli JA, Javid K, Cotrim K, Canabarro A, Louro RS, et al. Comparing the long-term success rates of tooth preservation and dental implants: a critical review. *J Funct Biomater* [Internet]. 2023; 14(3): 142. Available from: <https://doi.org/10.3390/jfb14030142>
10. Atieh MA, Almutairi Z, Amir-Rad F, Koleilat M, Tawse-Smith A, Ma S, et al. A retrospective analysis of biological complications of dental implants. *Int J Dent* [Internet]. 2022; 2022(1): 1545748. Available from: <https://doi.org/10.1155/2022/1545748>